

## Listing of claims:

For the convenience of the Examiner, all claims being examined are presented below.

1. **(Withdrawn)** A transgenic knockout mouse whose genome comprises a homozygous disruption in its endogenous FcRn gene, wherein said homozygous disruption prevents the expression of a functional FcRn protein, resulting in a transgenic knockout mouse in which exogenously administered IgG1 exhibits a substantially shorter half-life, as compared to the half-life of exogenously administered IgG1 in a wild-type mouse.
2. **(Withdrawn)** A transgenic knockout mouse whose genome comprises a homozygous disruption in its endogenous FcRn gene, wherein said homozygous disruption prevents the expression of a functional FcRn protein, resulting in a transgenic knockout mouse which is unable to absorb maternal IgG in the prenatal or neonatal stage of development.
3. **(Withdrawn)** The transgenic knockout mouse of Claim 1 or 2 wherein the homozygous disruption results from deletion or mutation of portions of the endogenous FcRn gene whereby a non-functional gene product or complete absence of the gene product is produced.
4. **(Withdrawn)** The transgenic knockout mouse of Claim 3 wherein the deleted portion of the endogenous FcRn gene is replaced with a neomycin-resistance cassette.
5. **(Withdrawn)** The transgenic knockout mouse of Claim 3 wherein the homozygous disruption consists of a deletion of 853 nucleotides of sequences upstream of exon 1, the entirety of exon 1 and intron 1, and 207 nucleotides immediately downstream of intron 2 of the FcRn gene, and wherein the deleted portion of the endogenous FcRn gene is replaced with a neomycin-resistance cassette.
6. **(Withdrawn)** A method of producing a monoclonal antibody which specifically binds FcRn through its antigen binding regions, comprising:
  - a) providing a FcRn- / - transgenic knockout mouse described in Claim 1;
  - b) immunizing the FcRn- / - transgenic knockout mouse with FcRn;

- c) generating a hybridoma which secretes a monoclonal antibody which specifically binds FcRn through its antigen binding regions from B-cells isolated from the immunized mouse; and
  - d) isolating the monoclonal antibody which specifically binds FcRn through its antigen binding regions from the hybridoma.
7. **(Withdrawn)** The monoclonal antibody produced by the method of Claim 6.
  8. **(Withdrawn)** The monoclonal antibody of Claim 7 which binds to human FcRn.
  9. **(Withdrawn)** The monoclonal antibody of Claim 7 which, when bound to FcRn, inhibits binding of FcRn to the Fc region of IgG.
  10. **(Withdrawn)** A monoclonal antibody which specifically binds FcRn through its antigen binding regions and, when bound to FcRn, inhibits binding of FcRn to the Fc region of IgG.
  11. **(Withdrawn)** The monoclonal antibody of Claim 7 which binds to the FcRn/ $\beta$ 2M complex but does not bind to  $\beta$ 2M alone or  $\beta$ 2M complexed with any other molecule other than FcRn and inhibits binding of the FcRn/ $\beta$ 2M complex to the Fc region of IgG.
  12. **(Withdrawn)** The monoclonal antibody of Claim 7 which inhibits the formation of the FcRn/ $\beta$ 2M complex by inhibiting binding of FcRn to  $\beta$ 2M.
  13. **(Withdrawn)** A method to detect FcRn molecules using the monoclonal antibody of Claim 7, said method consisting, but not limited to, enzyme-linked immunoassays (ELISA), radioimmunosorbent assay (RIA), immunoperoxidase-based in situ staining assay, antibody-based lateral flow assay and flow-cytometry assay.
  14. **(Withdrawn)** A mouse model for autoimmune disease in humans for use in developing human FcRn based therapeutics, the mouse being a transgenic knockout mouse whose genome comprises a homozygous disruption in its endogenous FcRn gene, wherein said homozygous disruption prevents the expression of a functional FcRn protein, the mouse also being transgenic for expression of functional huFcRn protein from a complete huFcRn gene.
  15. **(Withdrawn)** The mouse model of claim 14 wherein the transgenic knockout mouse is further characterized as having an NZM2410/J, BXSB/MpJ, BXSB/MpJ-

Yaa, MRL-MPJ, or MRL-MPJ-Faslpr background or any other mouse genotype backgrounds that are used in autoimmune studies.

16. **(Withdrawn)** The mouse model of claim 14 wherein the transgenic knockout mouse is further characterized as having the other known genes expressed in the muFcRn<sup>-/-</sup>, +huFcRn transgenic mouse that are important in autoimmune disease pathways.
17. **(Withdrawn)** The mouse model of Claim 14 wherein the huFcRn gene further comprises about 10 kb 5' and about 10 kb 3' sequences which flank the huFcRn gene in the human genome, and includes all known normal regulatory sequences.
18. **(Withdrawn)** A mouse model for autoimmune disease in humans for use in developing human FcRn based therapeutics, the mouse being a transgenic knockout mouse whose genome comprises a homozygous disruption in its endogenous FcRn gene, wherein said homozygous disruption prevents the expression of a functional FcRn protein, the mouse also being transgenic for expression of functional huFcRn protein from a huFcRn cDNA construct.
19. **(Withdrawn)** The mouse model of claim 18 wherein the transgenic knockout mouse is further characterized as having an NZM2410/J, BXSB/MpJ, BXSB/MpJ-Yaa, MRL-MPJ, or MRL-MPJ-Faslpr background or any other mouse genotype backgrounds that are used in autoimmune studies.
20. **(Withdrawn)** The mouse model of claim 18 wherein the transgenic knockout mouse is further characterized as having other known genes expressed in the muFcRn<sup>-/-</sup>, +huFcRn transgenic mouse that are important in autoimmune disease pathways.
21. **(Withdrawn)** The mouse model of Claim 18 wherein the huFcRn gene further comprises an FcRn cDNA construct containing regulatory sequences that produce variable to abnormally higher levels of protein expression than found with transgenes driven by endogenous FcRn regulatory sequences.
22. **(Withdrawn)** A mouse model for autoimmune disease in humans for use in developing human FcRn based therapeutics, the mouse being a transgenic knockout mouse whose genome comprises a homozygous disruption in its endogenous FcRn and IgG gene, wherein said homozygous disruption prevents the

expression of a functional FcRn and IgG protein, the mouse also being transgenic for expression of functional huFcRn protein from a huFcRn DNA construct and human IgG from a huIgG DNA construct, wherein the huFcRn protein in the mouse functions to protect human IgG from degradation.

23. **(Withdrawn)** A mouse model for autoimmune disease in humans for use in developing human FcRn based therapeutics, the mouse being a transgenic knockout mouse whose genome comprises a homozygous disruption in its endogenous FcRn gene, wherein said homozygous disruption prevents the expression of a functional FcRn protein, the mouse also being transgenic for expression of functional huFcRn protein from a huFcRn DNA construct and human IgG from a huIgG DNA construct, wherein the huFcRn protein in the mouse functions to protect human IgG from degradation.
24. **(Withdrawn)** A method to identify or characterize an inhibitor of FcRn mediated protection of IgG antibodies, comprising:
- a) providing functional huFcRn/hu $\beta$ 2M complexes in solution or bound to a support;
  - b) contacting candidate inhibitor to the huFcRn/hu $\beta$ 2M complex;
  - c) contacting human IgG to the huFcRn/hu $\beta$ 2M complex under conditions appropriate for binding of the huFcRn/hu $\beta$ 2M complex to the human IgG; and
  - d) assaying for inhibition of binding due to the candidate inhibitor, as compared to binding of a control huFcRn/hu $\beta$ 2M complex to human IgG in the absence of candidate inhibitor.
25. **(Withdrawn)** A method to identify or characterize an inhibitor of FcRn mediated protection of IgG antibodies, comprising:
- a) providing functional huFcRn/hu $\beta$ 2M complexes in solution or bound to a support;
  - b) contacting human IgG to the huFcRn/hu $\beta$ 2M complex under conditions appropriate for binding of the huFcRn/hu $\beta$ 2M complex to the human IgG;
  - c) contacting candidate inhibitor to the huFcRn/hu $\beta$ 2M; and

- d) assaying for inhibition of binding due to the candidate inhibitor, as compared to binding of a control huFcRn/hu $\beta$ 2M complex to human IgG in the absence of candidate inhibitor.
26. **(Withdrawn)** A method to identify or characterize an inhibitor of FcRn comprising:
- a) providing mammalian cells in culture which functionally express huFcRn;
  - b) contacting candidate inhibitor to the mammalian cells of step a);
  - c) contacting human IgG to the mammalian cells; and
  - d) assaying IgG catabolism by the mammalian cells in the presence and absence of candidate inhibitor, with an increase in IgG catabolism in the presence of candidate inhibitor, as compared to IgG catabolism in the absence of candidate being an indication that the candidate inhibits huFcRn.
27. **(Withdrawn)** A method to identify or characterize an inhibitor of FcRn comprising:
- a) providing mouse FcRn<sup>-/-</sup>,+huFcRn cells in culture;
  - b) contacting candidate inhibitor to the FcRn<sup>-/-</sup>,+huFcRn cells of step a);
  - c) contacting human IgG to the FcRn<sup>-/-</sup>,+huFcRn cells; and
  - d) assaying IgG catabolism by the mammalian cells in the presence and absence of candidate inhibitor, with an increase in IgG catabolism in the presence of candidate inhibitor, as compared to IgG catabolism in the absence of candidate inhibitor, being an indication that the candidate inhibitor inhibits huFcRn.
28. **(Withdrawn)** A method to identify or characterize an inhibitor of FcRn comprising:
- a) providing mammalian cells in culture which functionally express huFcRn;
  - b) contacting human IgG to the mammalian cells of step a);
  - c) contacting candidate inhibitor to the mammalian cells; and
  - d) assaying IgG catabolism by the mammalian cells in the presence and absence of candidate inhibitor, with an increase in IgG catabolism in the presence of candidate inhibitor, as compared to IgG catabolism in the

absence of candidate inhibitor, being an indication that the candidate inhibitor inhibits huFcRn.

29. **(Withdrawn)** A method to identify or characterize an inhibitor of FcRn comprising:
- a) providing mouse FcRn<sup>-/-</sup>,+huFcRn cells in culture;
  - b) contacting human IgG to the FcRn<sup>-/-</sup>,+huFcRn cells of step a);
  - c) contacting candidate inhibitor to the FcRn<sup>-/-</sup>,+huFcRn cells; and
  - d) assaying IgG catabolism by the mammalian cells in the presence and absence of candidate inhibitor, with an increase in IgG catabolism in the presence of candidate inhibitor, as compared to IgG catabolism in the absence of candidate inhibitor, being an indication that the candidate inhibitor inhibits huFcRn.
30. **(Currently Amended)** A method to identify or characterize an inhibitor of IgG protection by human FcRn (huFcRn) comprising:
- a) providing a first and a second transgenic mouse whose genome comprises a homozygous disruption in its endogenous FcRn gene, and whose genome further comprises a DNA sequence encoding a human FcRn operably linked to a regulatory sequence, wherein said homozygous disruption prevents the expression of a functional endogenous FcRn protein, and wherein the mouse expresses a functional human FcRn protein (a muFcRn<sup>-/-</sup>, +huFcRn transgenic mouse);
  - b) administering intravenously or intraperitoneally tracer human IgG and tracer human IgA to the first and the second transgenic mouse;
  - c) administering intravenously or intraperitoneally a candidate inhibitor to the first transgenic mouse; and
  - d) determining the half-life of the tracer human IgG and the half-life of the tracer human IgA in the first transgenic mouse which has received the candidate inhibitor and in the second transgenic mouse which has not received the candidate inhibitor, with a decrease in the half-life of the tracer human IgG, but not the tracer human IgA, in the first transgenic mouse, as compared to the half life of the tracer human IgG and the tracer human IgA

respectively in the second transgenic mouse, being an indication that the candidate inhibitor inhibits FcRn protection of IgG.

31. **(Withdrawn)** A method to inhibit FcRn mediated IgG protection in an individual comprising:
- a) providing a monoclonal antibody which specifically binds huFcRn through its antigen binding regions and inhibits binding of huFcRn to IgG; and
  - b) administering the monoclonal antibody of step a) to the individual in sufficient amounts to inhibit binding of huFcRn of the individual to IgG of the individual.
32. **(Withdrawn)** The method of Claim 31 wherein the individual has an autoimmune disease.
33. **(Withdrawn)** The method of Claim 31 wherein the individual has the autoimmune disease systemic lupus erythematosus.
34. **(Withdrawn)** The method of Claim 31 wherein the autoimmune disease is selected from the group consisting of, but not limited to, insulin resistant diabetes, myasthenia gravis, polyarteritis, cutaneous vasculitis, pemphigus vulgaris, Goodpasture's syndrome, rheumatoid arthritis, Kawasaki's disease, and Sjogren's syndrome.
35. **(Withdrawn)** A method to inhibit FcRn mediated IgG protection in an individual, comprising:
- a) providing an engineered molecule comprising an Fc region of IgG to which the FcRn binds; and
  - b) administering the engineered molecule of step a) to the individual in sufficient amounts to competitively inhibit binding of FcRn of the individual to IgG of the individual.
36. **(Withdrawn)** The method of Claim 35 wherein the individual has an autoimmune disease.
37. **(Withdrawn)** The method of Claim 35 wherein the autoimmune disease is lupus erythematosus.

38. **(Withdrawn)** The method of Claim 35 wherein the autoimmune disease is selected from the group consisting of, but not limited to, insulin resistant diabetes, myasthenia gravis, polyarteritis, cutaneous vasculitis, pemphigus vulgaris, goodpasture's syndrome, rheumatoid arthritis, Kawasaki's disease, and Sjogren's syndrome.
39. **(Withdrawn)** A method to inhibit FcRn mediated IgG protection in an individual, comprising:
- a) providing an organic molecule which binds to FcRn and inhibits binding of huFcRn to IgG; and
  - b) administering the organic molecule of step a) to the individual in sufficient amounts to inhibit binding of FcRn of the individual to IgG of the individual.
40. **(Withdrawn)** A method to inhibit FcRn mediated IgG protection in an individual, comprising:
- a) providing an inhibitor of FcRn expression to the individual; and
  - b) administering the inhibitor of step a) to the individual in sufficient amounts to inhibit expression of FcRn.
41. **(Withdrawn)** The method of Claim 40 wherein the inhibitor inhibits expression of FcRn at the transcription, translation, post-translational processing or protein transport to the membrane.
42. **(Withdrawn)** The method of Claim 40 wherein the inhibitor includes, but is not limited to, antisense oligonucleotides, reverse gene constructs, and specific targeting of regulator proteins.
43. **(Withdrawn)** A method to identify or characterize a therapeutic agent in the treatment of autoimmune diseases in humans, comprising:
- a) providing a mouse model of Claim 14;
  - b) administering a candidate therapeutic agent to the transgenic mouse of step a); and
  - c) monitoring autoimmune disease development in the transgenic mouse which receives the candidate therapeutic agent, with amelioration in autoimmune disease development, as compared to autoimmune disease development in a



transgenic mouse of step a) which does not receive the candidate therapeutic agent, being an indication that the candidate therapeutic agent may be useful for treating autoimmune diseases in humans.

44. **(Withdrawn)** A method to identify or characterize a therapeutic agent in the treatment of autoimmune diseases in humans, comprising:
- a) providing a mouse model of Claim 18;
  - b) administering a candidate therapeutic agent to the transgenic mouse of step a); and
  - c) monitoring autoimmune disease development in the transgenic mouse which receives the candidate therapeutic agent, with amelioration in autoimmune disease development, as compared to autoimmune disease development in a transgenic mouse of step a) which does not receive the candidate therapeutic agent, being an indication that the candidate therapeutic agent may be useful for treating autoimmune diseases in humans.
45. **(Withdrawn)** A method to identify or characterize a therapeutic agent in the treatment of autoimmune diseases in humans, comprising:
- a) providing a mouse model of Claim 22;
  - b) administering a candidate therapeutic agent to the transgenic mouse of step a); and
  - c) monitoring autoimmune disease development in the transgenic mouse which receives the candidate therapeutic agent, with amelioration in autoimmune disease development, as compared to autoimmune disease development in a transgenic mouse of step a) which does not receive the candidate therapeutic agent, being an indication that the candidate therapeutic agent may be useful for treating autoimmune diseases in humans.
46. **(Withdrawn)** A method to identify or characterize a therapeutic agent in the treatment of autoimmune diseases in humans, comprising:
- a) providing a mouse model of Claim 23;
  - b) administering a candidate therapeutic agent to the transgenic mouse of step a); and

- c) monitoring autoimmune disease development in the transgenic mouse which receives the candidate therapeutic agent, with amelioration in autoimmune disease development, as compared to autoimmune disease development in a transgenic mouse of step a) which does not receive the candidate therapeutic agent, being an indication that the candidate therapeutic agent may be useful for treating autoimmune diseases in humans.

**47-56. (Cancelled)**

57. **(Currently Amended)** A method to identify or characterize a candidate agent for FcRn-mediated drug stabilization, comprising:
- a) providing a formulation comprising a candidate agent for FcRn-mediated drug stability attached to a trackable composition;
  - b) administering intravenously or intraperitoneally the formulation to a muFcRn-/-, +huFcRn transgenic mouse, and also to a muFcRn-/- transgenic mouse; and
  - c) assaying the half-life of the formulation in the bloodstream of the muFcRn-/-, +huFcRn transgenic mouse and in the bloodstream of the muFcRn-/- transgenic mouse, with a substantially longer half-life of the formulation in the bloodstream of the muFcRn-/-, +huFcRn transgenic mouse than in the bloodstream of the muFcRn-/- transgenic mouse being an indication that the candidate agent promotes FcRn-mediated drug stabilization.
58. **(Previously Presented)** The method of Claim 57 wherein the candidate agent is derived from an IgG Fc region.
59. **(Previously Presented)** The method of Claim 57 wherein the candidate agent is derived from an IgG Fc region or fragment thereof, or other molecules which are structurally-similar or similar in sequence identity to the Fc-region of IgG.
60. **(Previously Presented)** The method of Claim 57 wherein the candidate agent is structurally-similar or similar in sequence identity to the Fc-region of IgG, such that the binding affinity of the candidate agent to the FcRn protein is greater than the binding affinity of the Fc-region of IgG to the FcRn protein.

61. **(Currently Amended)** A method for determining the difference in pharmacokinetics of a drug conferred by an FcRn-mediated drug stability candidate agent linked to the drug, comprising:
- a) providing a first trackable formulation comprising a drug linked to a FcRn-mediated drug stability candidate agent, and a second trackable formulation comprising the drug without the FcRn-mediated drug stability candidate agent;
  - b) administering intravenously or intraperitoneally the first formulation to a first muFcRn-/-, +huFcRn transgenic mouse, and the second formulation to a second muFcRn-/-, +huFcRn transgenic mouse;
  - c) determining the half-lives of the first and the second formulation in the bloodstream of the first and the second muFcRn-/-, +huFcRn transgenic mouse respectively; and
  - d) comparing the half-lives determined in step c) to one another, thereby determining the difference in half-lives conferred to the drug by the FcRn-mediated drug stability candidate agent.
62. **(Previously Presented)** The method of Claim 61 wherein the candidate agent is derived from an IgG Fc region.
63. **(Previously Presented)** The method of Claim 61 wherein the candidate agent is derived from an IgG Fc region or fragment thereof, or other molecules which are structurally-similar or similar in sequence identity to the Fc-region of IgG.
64. **(Previously Presented)** The method of Claim 61 wherein the candidate agent is structurally-similar or similar in sequence identity to the Fc-region of IgG, such that the binding affinity of the candidate agent to the FcRn protein is greater than the binding affinity of the Fc-region of IgG to the FcRn protein.
65. **(Withdrawn)** A method to identify or characterize a candidate agent for FcRn-mediated drug stabilization in a high-throughput system, comprising:
- a) providing a formulation comprising a candidate agent for FcRn-mediated drug stabilization attached to a trackable composition;

- b) administering the formulation to cells derived from muFcRn-/-, +huFcRn and muFcRn-/- or other mammalian cell lines which do not express FcRn; and
  - c) assaying the half-life of the formulation in cell media from cells in b), with a substantially longer half-life in the media from the muFcRn-/-, +huFcRn cells being an indication that the candidate agent promotes FcRn-mediated drug stabilization.
66. **(Withdrawn)** The method of Claim 65 wherein the candidate agent is derived from an immunoglobulin Fc region.
67. **(Withdrawn)** The method of Claim 65 wherein the candidate agent is derived from an Fc region or fragment thereof, or other molecules which are structurally-similar or similar in sequence identity to the Fc-region of IgG.
68. **(Withdrawn)** The method of Claim 65 wherein the candidate agent is structurally-similar or similar in sequence identity to the Fc-region of IgG, such that the engineered molecule binds with greater affinity to the FcRn protein.
69. **(Withdrawn)** A method to identify or characterize a candidate agent for FcRn-mediated drug stabilization in a high-throughput system, comprising:
- a) providing a formulation comprising a candidate agent for FcRn-mediated drug stabilization attached to a trackable composition;
  - b) administering the formulation to mammalian cells expressing huFcRn mammalian cell lines and mammalian cell lines which do not express FcRn; and
  - c) assaying the half-life of the formulation in cell media from cells in b), with a substantially longer half-life in the media from the mammalian cells expressing huFcRn being an indication that the candidate agent promotes FcRn-mediated drug stabilization.
70. **(Withdrawn)** The method of Claim 69 wherein the candidate agent is derived from an immunoglobulin Fc region.
71. **(Withdrawn)** The method of Claim 69 wherein the candidate agent is derived from an Fc region or fragment thereof, or other molecules which are structurally-similar or similar in sequence identity to the Fc-region of IgG.

72. **(Withdrawn)** The method of Claim 69 wherein the candidate agent is structurally-similar or similar in sequence identity to the Fc-region of IgG, such that the engineered molecule binds with greater affinity to the FcRn protein.
73. **(Withdrawn)** A method for determining the pharmacokinetics of an agent linked to an FcRn-mediated drug stability candidate, comprising:
- a) providing a trackable formulation comprising an agent linked to a candidate agent for FcRn-mediated drug stability;
  - b) administering the formulation to cells derived from muFcRn-/-, +huFcRn and muFcRn-/- or other mammalian cell lines which do not express FcRn; and
  - c) assaying the half-life of the formulation in cell media from cells in b), with a substantially longer half-life in the media from the muFcRn-/-, +huFcRn cells being an indication that the candidate agent promotes FcRn-mediated drug stabilization.
74. **(Withdrawn)** The method of Claim 73 wherein the candidate agent is derived from an immunoglobulin Fc region.
75. **(Withdrawn)** The method of Claim 73 wherein the candidate agent is derived from an Fc region or fragment thereof, or other molecules which are structurally-similar or similar in sequence identity to the Fc-region of IgG.
76. **(Withdrawn)** The method of Claim 73 wherein the candidate agent is structurally-similar or similar in sequence identity to the Fc-region of IgG, such that the engineered molecule binds with greater affinity to the FcRn protein.
77. **(Withdrawn)** A method for determining the pharmacokinetics of an agent linked to an FcRn-mediated drug stability candidate, comprising:
- a) providing a trackable formulation comprising an agent linked to a candidate agent for FcRn-mediated drug stability;
  - b) administering the formulation to mammalian cells expressing huFcRn mammalian cell lines and mammalian cell lines which do not express FcRn; and
  - c) assaying the half-life of the formulation in cell media from cells in b), with a substantially longer half-life in the media from the mammalian cells

expressing huFcRn being an indication that the candidate agent promotes FcRn-mediated drug stabilization.

78. **(Withdrawn)** The method of Claim 77 wherein the candidate agent is derived from an immunoglobulin Fc region.
79. **(Withdrawn)** The method of Claim 77 wherein the candidate agent is derived from an Fc region or fragment thereof, or other molecules which are structurally-similar or similar in sequence identity to the Fc-region of IgG.
80. **(Withdrawn)** The method of Claim 77 wherein the candidate agent is structurally-similar or similar in sequence identity to the Fc-region of IgG, such that the engineered molecule binds with greater affinity to the FcRn protein.
- 81-84. **(Cancelled)**
85. **(Currently Amended)** A method to identify or characterize a candidate agent for FcRn-mediated drug stabilization, comprising:
- a) providing a formulation comprising a candidate agent for FcRn-mediated drug stability formulated with a trackable composition;
  - b) administering intravenously or intraperitoneally the formulation to a muFcRn-/-, +huFcRn transgenic mouse, and also to a muFcRn-/- transgenic mouse; and
  - c) assaying the half-life of the formulation in the bloodstream of the muFcRn-/-, +huFcRn transgenic mouse and the bloodstream of the muFcRn-/- transgenic mouse, with a substantially longer half-life of the formulation in the bloodstream of the muFcRn-/-, +huFcRn transgenic mouse than in the bloodstream of the muFcRn-/- transgenic mouse being an indication that the candidate agent promotes FcRn-mediated drug stabilization.
86. **(Currently Amended)** A method for determining the difference in pharmacokinetics of a drug conferred by an FcRn-mediated drug stability candidate agent linked to the drug, comprising:
- a) providing a first trackable formulation comprising a drug linked to a FcRn-mediated drug stability candidate agent, and a second trackable formulation comprising the drug without the FcRn-mediated drug stability candidate agent;

- b) administering intravenously or intraperitoneally the first and the second formulation to a muFcRn<sup>-/-</sup>, +huFcRn transgenic mouse;
- c) determining the half-lives of the first and the second formulation in the bloodstream of the muFcRn<sup>-/-</sup>, +huFcRn transgenic mouse; and
- e) comparing the half-lives determined in step c) to one another, thereby determining the difference in half-lives conferred to the drug by the FcRn-mediated drug stability candidate agent.